



Selective Isotope-Labeling Methods for Protein Structural Studies

Hanudatta S. Atreya, PhD

NMR Research Centre, Indian Institute of Science, Bangalore, India

One of the major contributing factors to the rapid advance of biomolecular NMR spectroscopy is the emergence of different isotope labeling methods. Recent developments in biotechnology have made it easier and economical to introduce ^{13}C , ^{15}N and ^2H into proteins and nucleic acids. At the same time, there has been an explosion in the number of NMR experiments that utilize such isotope-labeled samples. Thus, a combination of isotopic labeling and multidimensional, multinuclear experiments has significantly expanded the range of problems in structural biology amenable to NMR.

Isotope labeling in proteins can be broadly classified into four categories: Uniform, amino acid-type selective, site-specific and random/fractional labeling. The beginning of systematized isotope labeling in proteins can be traced back to late 60's in the group of Jardetsky and Katz and coworkers.^{1,2} Theirs was also one of the first amino acid-type selective labeling methods involving incorporation of specific protonated amino acids against a deuterated background. In the 80's uniform ($^{13}\text{C}/^{15}\text{N}$) and selective incorporation of ^{15}N -labeled amino acids against an unlabeled ($^{12}\text{C}/^{14}\text{N}$) background was developed.³ Subsequently, a variety of labeling methods have emerged (reviewed in [4] and [5] and illustrated in Figure 1).

In addition to uniform ($^{13}\text{C}/^{15}\text{N}/^2\text{H}$) labeling, amino acid-type or site-selective labeling is often pursued as it helps in spectral simplification and provides specific probes for structural and

dynamic studies. Selective amino acid-type labeling also aids in sequence-specific resonance assignments by helping to identify resonances which are otherwise buried in the crowded regions of 2D and 3D NMR spectra. However, a disadvantage of this method is the possible mis-incorporation of ^{15}N label in undesired amino acids (also called as "isotope scrambling").³ This happens due to metabolic conversion of one amino acid to another in the bio-synthetic pathway of the cell. The problem becomes more severe for amino acids higher up or intermediates in the metabolic pathway such as Asp, Glu and Gln (See Figure 2 showing the biosynthetic pathway in *E. coli*). For those which are end-products in the production pipeline (Ala, Arg, Asn, Cys, His, Ile, Lys, Met, Pro and Trp) isotope scrambling is minimal and the remaining (Gly, Phe, Leu, Ser, Thr, Tyr and Val) have medium to weak interconversion. Isotope scrambling in *E. coli* can be minimized by reducing the activity of the enzyme(s) catalyzing the inter-conversion or amino transfer using either specific (auxotrophic) strains³ or using enzyme inhibitors.⁶ Another alternative is to use cell-free or *in vitro* expression systems which lack these enzymes.⁴

One drawback of amino acid selective labeling is the expense associated with the use of $^{13}\text{C}/^{15}\text{N}$ labeled amino acids. A relatively inexpensive method is that of amino acid selective "unlabeling" or reverse labeling. In this method, the host organism is grown on a medium containing the desired unlabeled (*i.e.*, $^1\text{H}/^{12}\text{C}/^{14}\text{N}$) amino acid against a labeled ($^{13}\text{C}/^{15}\text{N}$) background. This is somewhat akin to the selective protonation experiment by Jardetsky¹ and Katz.² Reverse labeling was first used by Bax and coworkers⁷ and developed further by other groups for different applications.^{8,9,10} The problem of isotope scrambling (in this case being the mis-incorporation of ^{14}N) remains largely the same as in the selective-labeling approach mentioned above (for a detailed table of possible scrambling of ^{14}N see reference 10).

In addition to the above, new isotope-labeling methods continue to be developed. More recent methods of segmental labeling¹¹ and stereo-arrayed isotope labeling (SAIL)¹² open up new avenues in protein structural studies. The future points towards a combination of different isotope-labeling methods to address challenging and complex problems in structural biology.




Figure 1. Different isotope-labeling methods.

(continued)

References

1. Markley, J.K.; Putter, I.; and Jardetsky, O. **1968**. High-resolution nuclear magnetic resonance spectra of selectively deuterated staphylococcal nuclease. *Science*, *161*, 1249-1251.
2. Crespi, H.L.; Rosenberg, R.M.; and Katz, J.L. (1968) **1968**. Proton magnetic resonance of proteins fully deuterated except for H-leucine side chains. *Science*, *161*, 795-796.
3. Muchmore D.D.; McIntosh, L.P.; Russell, C.B.; Anderson, D.E.; Dahlquist, F.W. **1989**. Expression and nitrogen-15 labeling of proteins for proton and nitrogen-15 nuclear magnetic resonance. *Methods Enzymol*, *177*, 44-73.
4. Ohki, S. and Kainosho, M. **2008**. Stable isotope labeling for protein NMR. *Prog NMR Spectrosc*, *53*, 208-226.
5. *J. Biomol NMR*, **2010**. Special issue: Vol 46: 1-125.
6. Tong, K.I.; Yamamoto, M.; and Tanaka, T. **2008**. A simple method for amino acid selective isotope labeling of recombinant proteins in *E. coli*. *J Biomol NMR*, *42*, 59-67.
7. Vuister, G.W.; Kim, S.J.; Wu, C., Bax, A. **1994**. 2D and 3D NMR study of phenylalanine residues in proteins by reverse isotopic labeling. *J Am Chem Soc*, *116*, 9206-9210.
8. Shortle, D. **1995**. Assignment of amino acid type in ^1H - ^{15}N correlation spectra by labeling with ^{14}N amino acids. *J Magn Reson B*, *105*, 88-90.
9. Atreya, H.S. and Chary, K.V.R. **2001**. Selective unlabeled of amino acids in fractionally ^{13}C -labeled proteins: An approach for stereospecific NMR assignments of CH₃ groups in Val and Leu residues. *J Biomol NMR*, *19*, 267-272.
10. Krishnarjuna, B.; Jaiupuria, G.; Thakur, A.D.; Silva, P.; and Atreya, H.S. **2010**. Amino acid selective unlabeled for sequence specific resonance assignments in proteins. *J Biomol NMR*, *49*, 38-51.
11. Cowburn, D. and Muir, T.W. **2001**. Segmental isotopic labeling for structural biological applications of NMR. *Methods Enzymol*, *339*, 41-54.
12. Kainosho, M.; Torizawa, T.; Iwashita, Y.; Terauchi, T.; Ono, A.M.; and Peter Güntert. **2006**. Structure of the putative 32 kDa myrosinase binding protein from arabidopsis (At3g16450.1) determined by SAIL-NMR. *Nature*, *440*, 52-57.